

Entada rheedii Sprengel, Sprengel

Identifiants : 12743/entrhe

Association du Potager de mes/nos Rêves (<https://lepotager-demesreves.fr>)

Fiche réalisée par Patrick Le Ménahèze

Dernière modification le 12/05/2024

- Classification phylogénétique :

- Clade : Angiospermes ;
- Clade : Dicotylédones vraies ;
- Clade : Rosidées ;
- Clade : Fabidées ;
- Ordre : Fabales ;
- Famille : Fabaceae ;

- Classification/taxinomie traditionnelle :

- Règne : Plantae ;
- Division : Magnoliophyta ;
- Classe : Magnoliopsida ;
- Ordre : Fabales ;
- Famille : Fabaceae ;
- Genre : Entada ;

- Synonymes : *Adenantha gogo* Blanco, *Entada gigas* G. Gilbert & Boutique, *Entada gogo* (Blanco) I. M. Johnston, *Entada monostachya* DC, *Entada pursaetha* DC, *Entada rheedii* subsp. *rheedii*, *Entada schefferi* Ridl, *Entada cirrhosa* Raf, ;

- Nom(s) anglais, local(aux) et/ou international(aux) : Sea bean, Mackay bean, , Angkunh, Bendo, Bolcchak chhan, Bumburandje, Cariu, Elephant climber, Gila, Gumpe, Gunguro, Irikki, Kaka valli, Kakkankai, Kawi-hrui, Makkinkokka, Malam thellukka, Manga kodi, Mufwihi, Pangra, Sanguzi, Vor angkhournh, Wijang, Yanaikozhinji, Zangusi ;



- Rapport de consommation et comestibilité/consommabilité inférée (partie(s) utilisable(s) et usage(s) alimentaire(s) correspondant(s)) :

Parties comestibles : feuilles, graines, légumes^{{}(0(+x)) (traduction automatique)} | Original : Leaves, Seeds, Vegetable^{{}(0(+x))} Les graines sont utilisées dans la bouillie comme aliment de famine. Ils sont cuits, pilés et trempés dans l'eau pendant une période prolongée, puis recuits. Les graines sont transformées en pâte avec de l'eau et du cumin. Attention: Les graines sont toxiques lorsqu'elles sont crues. Les jeunes feuilles sont consommées comme légume. Les jeunes feuilles tendres sont bouillies et mangées



cf. consommation

- Illustration(s) (photographie(s) et/ou dessin(s)):

- Liens, sources et/ou références :

dont classification :

dont livres et bases de données : ⁰"Food Plants International" (en anglais) ;

dont biographie/références de ⁰"FOOD PLANTS INTERNATIONAL" :

Arinathan, V., et al, 2007, Wild edibles used by Palliyars of the western Ghats, Tamil Nadu. *Indian Journal of Traditional Knowledge*. 6(1) pp 163-168 ; Binu, S., 2010, Wild edible plants by the tribals in Pathanamthitta district, Kerala. *Indian Journal of Traditional Knowledge*. 9(2): 309-312 ; Bruschi, P., et al, 2014, Traditional use of plants in a rural community of Mozambique and possible links with Miombo degradation and harvesting sustainability. *Journal of Ethnobiology and Ethnomedicine*. 2014, 10:59 ; Burkill, H. M., 1985, The useful plants of west tropical Africa, Vol. 3. Kew. ; Cengel, D. J. & Dany, C., (Eds), 2016, Integrating Forest Biodiversity Resource Management and Sustainable Community Livelihood Development in the Preah Vihear Protected Forest. *International Tropical Timber Organization* p 123 (As Entada pursaetha) ; Chapman, J. D. & Chapman, H. M., 2001, The Forest Flora of Taraba and Andamawa States, Nigeria. *WWF & University of Canterbury*. p 184 ; Dunlop, C.R., Leach, G.J. & Cowie, I.D., 1995, Flora of the Darwin Region. Vol 2. Northern Territory Botanical Bulletin No 20. p 23 (As Entada pursaetha) ; Grubben, G. J. H. and Denton, O. A. (eds), 2004, Plant Resources of Tropical Africa 2. Vegetables. *PROTA*, Wageningen, Netherlands. p 561 ; ILDIS Legumes of the World <http://www.ildis.org/Legume/Web> ; Kar, A., et al, 2013, Wild Edible Plant Resources used by the Mizos of Mizoram, India. *Kathmandu University Journal of Science, Engineering and Technology*. Vol. 9, No. 1, July, 2013, 106-126 (As Entada pursaetha) ; Lungphi, P., Wangpan, T. & Tangjang, S., 2018, Wild edible plants and their additional uses by the Tangsa community living in the Changlang district of Arunachal Pradesh, India. *Pleione* 12(2): 151 - 164. 2018. ; Phon, P., 2000, Plants used in Cambodia. Â© Pauline Dy Phon, Phnom Penh, Cambodia. p 275 (As Entada pursaetha) ; Prodr. 2:425. 1825 (As Entada pursaetha) ; Ramachandran, V.S., 1987, Further Notes on the Ethnobotany of Cannanore District, Kerala. *J. Econ. Tax. Bot.* Vol. 11 No. 1 pp 47- (As Entada pursaetha) ; Reitseveld, S., 2013, The Animals and Plants of the Zazamalala Forest in Western Madagascar. p 78 ; Royal Botanic Gardens, Kew (1999). Survey of Economic Plants for Arid and Semi-Arid Lands (SEPASAL) database. Published on the Internet; <http://www.rbghkew.org.uk/ceb/sepasal/internet> [Accessed 28th April 2011] ; Singh, B., et al, 2012, Wild edible plants used by Garo tribes of Nokrek Biosphere Reserve in Meghalaya, India. *Indian Journal of Traditional Knowledge*. 11(1) pp 166-171 ; Sukarya, D. G., (Ed.) 2013, 3,500 Plant Species of the Botanic Gardens of Indonesia. *LIPI* p 923 ; Sundriyal, M., et al, 2004, Dietary Use of Wild Plant Resources in the Sikkim Himalaya, India. *Economic Botany* 58(4) pp 626-638 (As Entada pursaetha) ; Verdcourt, B., 1979, Manual of New Guinea Legumes. *Botany Bulletin* No 11, Division of Botany, Lae, Papua New Guinea. p 135 (As Entada pursaetha) ; White, F., Dowsett-Lemaire, F. and Chapman, J. D., 2001, Evergreen Forest Flora of Malawi. Kew. p 316